

DIAGNOSTICS OF PREGNANT WOMEN WITH GESTATIONAL DIABETES

Karimova Gulrukh Komilovna

Department of Obstetrics and Gynecology №1, Bukhara State Medical Institute. Abu Ali Ibn Sina of the Ministry of Health of Uzbekistan.

DOI: 10.47750/pnr.2022.13.508.408

Abstract

Object: Selection of an effective method of delivery for mother and child in pregnant women with gestational diabetes.

Purpose of the study: Selecting the optimal method of delivery by studying the outcome of labor for the mother and fetus in women with gestational diabetes.

Material and methods: The retrospective group included 67 women who passed through the regional perinatal center, city maternity complex and family clinics No. 5, No. 6 of the city of Bukhara in the periods from 2016 to 2018. The main (prospect) group consisted of 68 women whose pregnancy proceeded against the background of overt or gestational diabetes mellitus (GDM). The control group consisted of 31 women whose pregnancy proceeded physiologically.

Results and discussion: As a result of the study, the course of pregnancy and childbirth according to the history of childbirth and own observations revealed that in all (60%) pregnant women with GDM, pregnancy and childbirth proceeded with any complications, like in the mother and the fetus.

Conclusion: Thus, the optimal delivery route is selected taking into account the condition of the fetus and the level of TSH and clay hemoglobin. If pregnancy proceeded against the background of corrected diabetes with the use of the optimal insulin regimen and rational diet therapy, the absence of signs of diabetic fetopathy, then in this category of pregnant women, the delivery through the birth canal is considered the best option. The issue of delivery time is decided individually, taking into account the degree of compensation for diabetes mellitus, cervical maturity, condition and size of the fetus.

Keywords: gestational diabetes mellitus, carbohydrate metabolism, delivery, preeclampsia.

Introduction

The problem of pregnancy and childbirth with gestational diabetes mellitus (GDM) is very relevant and is still not completely resolved. This is due to an increase in the number of pregnant women with this pathology, associated with a sharp increase in the incidence of diabetes in the population, as well as an improvement in the quality of diagnosis. The prevalence of all forms of diabetes mellitus among pregnant women reaches from 3.5% to 7.6%. Perinatal mortality in pregnancy complicated by diabetes mellitus of any type is 30-50%. After 3 months after giving birth, 4 out of 100 women with gestational diabetes develop a typical clinical picture of type 2 diabetes mellitus, after 1 year in 32% and after 8 years in 46% [14].

According to large-scale epidemiological studies, GDM is diagnosed in approximately 4% of pregnant Caucasian races. The prevalence of GDM can vary from 1 to 14% (an average of 7%), which depends on the analyzed population of women and the frequency of use of the oral glucose tolerance test (OGTT) used to diagnose the

disease. A screening and diagnostic program for GDS is also poorly organized. More over, according to WHO in the EU countries and the USA in 2009, 230,000 cases of GDM were recorded [2,8,11].

In most patients, gestational diabetes occurs with unexpressed, rare, and lack of clinical symptoms. These women tend to be older of reproductive age (35-49 years old) and they have found a bowl obesity [3, 11].

Distinctive features of pregnant women after the use of assisted reproductive technologies (ART) are age-related risk (more than 70% of patients over 30 years old); high incidence of multiple pregnancy; gynecological diseases and surgical intervention in the anamnesis; the presence of somatic pathology, hypertension of hypertension and hormonal disorders: obesity, polycystic ovary syndrome (PCOS). These pathological conditions significantly increase the risk of obstetric and perinatal complications [9 ,12].

The most significant risk factors for GDM according to data (T. Kovalenko 2007) are: age over 30 years (55.4%), obesity (49.2%), family history of diabetes burdened by diabetes (46.6%), glucosuria (43%) and fasting hyperglycemia (52%). It has been established that the presence of 2 or more factors increases the risk of GDM by 10 times or more.

The high frequency of multiple pregnancy with GDM may be caused almost double the concentration, and consequently, the double pin and rinsulyarnym action placental hormone, cortisol, etc. due to the functioning of several placentas [5,15].

The occurrence in pregnant women with gestational diabetes is facilitated by the high sensitivity of the fetus in the antenatal period to the action of any endogenous and exogenous damaging factors. F actors risk of developing GDM in modern obstetric diabetology can be divided into 2 groups [10].

According to the results of Gafurov M.R. (2014) the gestational age at which GDM was detected ranged from 21 to 34 weeks. An individual diet was selected for the patients, which was effective in 16 (57.1%), while 12 (42.9%) pregnant women were supplemented with insulin therapy. All patients performed glycemic self-control and kept a food diary. In addition to GDM, the course of pregnancy was complicated by preeclampsia in 4 (14.3%) women, in one case in the form of severe hepatosis , and in 50% of patients with the threat of abortion. Fetoplacental insufficiency (FPI) was observed in 4 (14.3%) patients with a single pregnancy and in all cases of multiple pregnancy [7,13,1,14].

Women who are at a constant subcutaneous infusion of insulin via insulin pump, during delivery of insulin administration is continued with the standard basal rate. After separation of the placenta, the infusion rate is reduced by 2 times and the intravenous infusion of the glucose-potassium mixture begins , complete cancellation of insulin is possible [4,10].

Women with GDM are at risk for developing type 2 diabetes after childbirth, therefore, they should be observed in the future prevention program for this disease. In women with a history of GDM, low-dose estrogen-progestogen oral contraceptives may be used if there are no medical contraindications. Children of mothers with GDM should be observed for the development of obesity or impaired glucose tolerance [6].

Thus, the organization and implementation of a universal program for screening and diagnosis of GDS, management of pregnancy and childbirth will improve pregnancy outcomes for both mother and future offspring and can be the basis for the prevention of type 2 diabetes in the future in this category of women and their children. Timely diagnosis and rational management of pregnancy in women with GDM, the use of medical methods and diet therapy can reduce the risk of complications during pregnancy and contribute to the birth of healthy children.

Purpose of the study

Selection of the optimal method rod or permitted by the study outcome childbirth for mother and fetus in women with gestational diabetes.

Material and methods

In the retrospective group included 67 women who passed through the regional perinatal center and number 5, number 6 family polyclinics of the city of Bukhara in the periods from 2016 to 2018 y. The main group consisted of 67 women whose pregnancy proceeded against the background of obvious or gestational diabetes. The control group consisted of 31 women whose pregnancy proceeded physiologically.

The work was performed on the basis of the Department of Obstetrics and Gynecology of the Medical institute and maternity hospitals in Bukhara. Gathering a diagnostic history through a survey, we specified the following data. Clinical diagnostic methods for women in the prospective group included: questioning 68 pregnant women and assessing the risk of developing GDH, consulting an endocrinologist, measuring blood pressure, body weight, calculating body mass index (BMI) using the Kettle formula: $BMI = \text{body weight before pregnancy, kg} / (\text{growth, m})^2$, general clinical tests of blood and urine, taking into account risk factors for the development of GDM. Ultrasound of the uterus, fetus in dynamics, glycated hemoglobin level, glucose tolerance test (TT to G) and other $BMI > 25 \text{ kg} / \text{m}^2$, presence of relatives of the 1st line of relationship with GDM or other history of carbohydrate metabolism, glycosuria during this pregnancy, age, women over 30 years, the birth of a child weighing more than 4000 g or a history of stillbirth, birth of children with congenital malformations in the history of habitual miscarriage pregnancy, a history of polyhydramnios, abnormal weight gain during this belt. Below are data on the age category of women in the examined groups.

Results and discussion

Some pregnancy complications, being characteristic, are not only for GDM, which are more often observed in women with this disease. When studying the course of pregnancy and childbirth according to the history of childbirth and own observations, it was found that in all (100%) pregnant women with GDM, pregnancy and childbirth proceeded with any complications. And in most cases, several complications were observed in combination, both in the mother and in the fetus. Hypertensive pathologies joined every second woman in a retrospective group.

Pregnancy was complicated by polyhydramnios in 21 cases in the main group, which amounted to 31%, 14 (24%) cases in the retrospective group, which was caused by infections of various nature. Basically, this complication was observed in women who underwent ARI during pregnancy.

In the control group, polyhydramnios was observed in 1 cases (9%), which was confirmed by clinical signs and ultrasound data. Observing the postpartum period in groups of women with GDM revealed that in 13 women (22.8% of cases) this period was complicated by early bleeding, which may have a mixed etiology (atonic and coagulopathy origin), postpartum hemorrhage is more often observed, which may be associated with overstretching uterus with polyhydramnios and large fetal size. Basically, bleeding was observed in the first 2 hours after separation and separation of the placenta and with a newborn's body weight of more than 4000 grams during prolonged labor. Macrosomia with a fetal weight of more than 4,500 grams, as you know, is the result of an excess supply of glucose to the fetus through the uteroplacental system.

In our cases, macrosomia occurred in 56% of cases in the main group and in 27.3% of cases in the control group (table number 1).

Table No. 1 Obstetric complications of gestational diabetes

№	Obstetric complications of GDM	Retrospective		Main		Control abs n = 11%	
		abs n = 67%	abs n = 100%	abs n = 68%	abs n = 65.4%	abs n = 0	abs n = 0
1	Preeclampsia (PE).	67	100	37	65.4	0	0

2	Urinary tract infections.	21	31.3	27	47.4	1	9
3	Premature rupture of the membranes	28	41.8	31	54.4	9	81.8
4	Macrosomia of the fetus.	67	100	32	56	2	18
5	Cesarean section	40	60	23	41	3	27.3
6	Polyhydramnios and infection.	21	31	14	24	1	9
7	Postpartum hemorrhage.	15	22	13	22.8	1	9

As you know, in newborns from mothers with diabetes, the risk of developing congenital anomalies increases by 3 times compared with 1-2% of the basic risk for all newborns. Most often, heart defects and abnormalities in the development of limbs occur. A typical, but rather rare anomaly is agenesis of the sacrum.

Complications of the postpartum period

The method of delivery and the course after in the comparative aspect are also studied. When studying the history of childbirth in women in a retrospective group, it was revealed that mainly delivery methods were operational. A caesarean section in this group of patients was performed in 68.6% of cases, childbirth flowed through the natural birth canal. A high percentage of operative births indicate that the pregnancy in these women occurred against the background of uncorrected hyperglycemia, therefore, with complications from the mother and fetus, which ultimately were indications for abdominal delivery (table number 1).

In almost all cases, childbirth was complicated by various complications. These were mainly: bleeding in excess of 500 ml during childbirth per vias naturals and more than 1000 ml during cesarean section. Hemorrhage, bearing atopic and coagulopathic character followed by 81% operational and 45.6% conservative. Given the high risk, two pairs of vessels were ligated (a. Ovaria propriety et a. Rotundum uteri). In 9 women, by examination, the operations were expanded to amputation of the uterus (5-23.8%) and extirpation (4-19%) of the uterus.

Maternal mortality was registered in 1 case, which amounted to 4.8%. the postpartum and postoperative period was also accompanied by complications, especially in women who underwent caesarean section (52.3%). These women have been observed sub involution uterus, accompanied lohiometrey and hematoma. In all likelihood, this was due to inhibition of the contractile ability of the uterus due to prolonged overstretching of its muscles and the phenomena of septic complications.

In the prospective group, 23 women had a cesarean section, they accounted for 34%. It was possible to reduce the operative birth mainly due to rational diet therapy and the choice of optimal individual insulin therapy.

Conclusions

Thus, and Learn the birth outcomes, monitor the progress of post-natal and post-operative period, revealed that pregnancy in 65.4% of women with GDM complicated by hypertensive disorders. In almost every third (31.6%) pregnant women, polyhydramnios of varying degrees joined. Delivery in (41%) cases ended with cesarean section, 45.6% of them were complicated by bleeding. Every second case from the retrospective and every 4 case from the prospective group was complicated by suture infiltration (50.5% and 25.3%, respectively).

The optimal delivery route is selected taking into account the condition of the fetus and the level of TSH and glycated hemoglobin. If pregnancy proceeded against the background of corrected diabetes with the use of the optimal insulin regimen and rational diet therapy, the absence of signs of diabetic fetopathy, then in this category of pregnant women, the delivery through the birth canal is considered the best option. The issue of delivery time is decided individually, taking into account the degree of compensation for diabetes mellitus, cervical maturity, condition and size of the fetus.

Bibliography:

1. Дерябина, Е.Г. Старая и новая концепции гестационного сахарного диабета: влияние на распространенность и перинатальные исходы / Е.Г.Дерябина // Сахарный диабет – пандемия XXI века : сборник тезисов VIII Всероссийского диабетологического Конгресса.- Москва, 2018 –С. 398.
2. Карапетян Т.Э. Генитальные микоплазмы и исходы беременности / Т.Э. Карапетян, В.В. Муравьева, А.С. Анкирская // Проблемы репродукции. - 2012. № 3. - С. 88-93.
3. Каримова Г.К., Наврузова Н.О. и Нуриллоева Ш.Н. (2020). Индивидуальный подход к ведению гестационного диабета. Европейский журнал молекулярной и клинической медицины , 7 (2), 6284-6291.
4. Karimova, G. K., Navruzova, N. O., & Nurilloeva Sh, N. (2020). An individual approach to the management of gestational diabetes. European Journal of Molecular & Clinical Medicine, 7(2), 6284-6291.
5. Karimova, G. K., Ikhtiyarova, G. A., & Muminova, N. K. (2021). Early biochemical markers and screening diagnosis of gestational diabetes mellitus and its prevention during pandemic period. Journal of Natural Remedies, 22(1 (1)), 17-26.
6. Каримова Г.К., Ихтиярова Г.А. и Муминова Н.К. (2021). Ранние биохимические маркеры и скрининг-диагностика гестационного сахарного диабета и его профилактика в период пандемии. Журнал природных средств правовой защиты , 22 (1 (1)), 17-26.
7. Sh, I. A., Ikhtiyarova, G. A., Musaeva, D. M., & Karimova, G. K. (2020). Assessment of the status of pregnant women with diabetes mellitus infected with COVID-19. New day in medicine, 2, 30.
8. Ш, И.А., Ихтиярова, Г.А., Мусаева, Д.М., Каримова, Г.К. (2020). Оценка состояния беременных с сахарным диабетом, инфицированных COVID-19. Новый день в медицине , 2 , 30.
9. Иноятов, А. Ш., Ихтиярова, Г. А., Мусаева, Д. М., & Каримова, Г. К. (2020). Оценка состояния беременных женщин с диабетом, при заражении COVID-19. Новый день в медицине, (2), 101-103.
10. Ихтиярова Г.А., Каримова Г.К., Наврузова Н.О. (2019). ХайруллаевЧ. К. Ультразвуковая диагностика диабетической фетопатии у беременных с метаболическим синдромом на фоне сахарного диабета. Медицина и спорт2019 , (3-4), 56-58.
11. Ikhtiyarova, G. A., Karimova, G. K., & Navruzova, N. O. (2019). KhairullaevCh. K. Ultrasound diagnostics of diabetic fetopathy in pregnant women with metabolic syndrome on the background of diabetes mellitus. Medicine and sports2019, (3-4), 56-58.
12. Каримова, Г. К., Ихтиярова, Г. А., & Наврузова, Н. О. (2020). Скрининг диагностика гестационного диабета. Новый день в медицине, (1), 220-222.
13. Ashurova, N. G., Karimova, G. K., & Norov, F. H. (2018). Modern views on a problem of the gestational diabetes. doctors newsletter, (1), 94-98.
14. Каримова, Г. К. (2022). Гестацион қандли диабетни эрта ташхислашнинг биокимёвий скрининги. Barqarorlik va yetakchi tadqiqotlar onlayn ilmiy jurnali, 2(8), 199-212.
15. Ихтиярова, Г. А., Каримова, Г. К., Наврузова, Н. О., & Хайруллаев, Ч. К. (2019). Ультразвуковая диагностика диабетической фетопатии у беременных с метаболическим синдромом на фоне сахарного диабета. Тиббиёт ва спорт, (3-4), 56-58.

16. Хамдамов И.Б., Хамдамов А.Б. и Каримова Г.К. (2016). Кахаров И.И. Гормональная контрацепция у женщин с сахарным диабетом. Молодежный инновационный вестник , (1), 209-210.
17. Khamdamov, I. B., Khamdamov, A. B., & Karimova, G. K. (2016). Kakharov II Hormonal contraception in women with diabetes. Youth Innovative Bulletin, (1), 209-210.
18. Комиловна, КГ . (2022). Современные взгляды на проблему гестационного сахарного диабета. Международный журнал систем здравоохранения и медицинских наук , 1 (4), 344–350.
19. Komilovna, K. G. . (2022). Modern Views on the Problem of Gestational Diabetes Mellitus. International journal of health systems and medical sciences, 1(4), 344–350.
20. Тошева, И., Ашурова, Н., & Ихтиярова, Г. (2020). Разрыв плодных оболочек в недоношенном сроке, как фактор развития акушерских осложнений. Журнал вестник врача, 1(1), 77-80.
21. 55. Тошева, И. И., & Ихтиярова, Г. А. (2020). Патоморфология последов, осложнения беременности, родов и исходы новорожденных с дородовым излитием околоплодных вод. Opinion leader, (2), 56-60.
22. 56. Тошева, И. И., & Ихтиярова, Г. А. (2020). Исходы беременности при преждевременном разрыве плодных оболочек. РМЖ. Мать и дитя, 3(1), 16-19.
23. 57. Тошева, И. И., & Ашурова, Н. Г. (2019). Исходы родов у беременных с преждевременным излитием околоплодных вод. Вестник Дагестанской государственной медицинской академии, (4), 34-37.
24. 58. Тошева, И. И., Ихтиярова, Г. А., & Аслонова, М. Ж. (1999). Современные методы индукции родов у женщин с отхождением околоплодных вод с внутриутробными инфекциями. Инфекция, иммунитет и фармакология, 254.
25. 59. Тошева И.И., Ашурова Н.Г., Ихтиярова Г.А. Разрыв плодных оболочек в недоношенном сроке, как фактор развития акушерских осложнений//Журнал Проблемы биологии и медицины. - 2020. - №1. - С.76-79.
26. 60. Тошева И.И., Ихтиярова Г.А. Дифференцированные подходы к методам родоразрешения при хориоамнионите. Вестник оперативной хирургии и топографической анатомии № 1 (01), Том 1, ISSN 2713–3273. 2020: 25–29.
27. 61. Тошева И.И., Каримова Г.К., Адизова С.Р. Изучение причин акушерских осложнений на фоне излитие околоплодных вод в доношенном сроке. Вестник Ташкентской медицинской академии. 2020:170-171.
28. Тошева И.И., Мусаходжаева Д.А., Магзумова Н.М. Родовозбуждение при антенатальной гибели плода у женщин с излитием околоплодных вод и внутриутробной инфекцией. Теоретической и клинической медицины Том 1, № 6 2021: 111–113.
29. Tosheva, I. I., Ikhtiyarova, G. A., & Aslonova, M. J. (2019). Introduction of childbirth in women with the discharge of amniotic fluid with intrauterine fetal death. Problems and solutions of advanced scientific research, 1(1), 417-424.
30. Tosheva, I. I., & Ikhtiyarova, G. A. (2019). Obstetric complications in pregnant women with premature discharge of amniotic fluid. Biologiya va tibbiy muammolari, 42(115), 146-149.
31. Tosheva II, Ikhtiyarova GA Cytokine Profile Changing in Pregnant Women with Chorioamnionitis// Open Access Journal of Gynecology-2021.6(4): 000227. P.1-6.
32. Tosheva Iroda Isroilovna. (2022). Labor and Perinatal Outcomes in Women with Preterm Rupture of Membranes and with Chorioamnionitis. INTERNATIONAL JOURNAL OF HEALTH SYSTEMS AND MEDICAL SCIENCES, 1(4), 297–305.
33. Isroilovna, T. I. . (2022). Preterm Rupture of Membranes, As a Factor in the Development of Obstetrics Complications. International journal of health systems and medical sciences, 1(4), 336–343.