

"Screening Of Juniperus Communis Linn. Oil For Anthelmintic, Injury Recuperating And Hepatoprotective Exercises In Rodents"

Lalit Bisht ¹, Ranjana Sonkar ², Arun Kumar Maurya ³

¹ Lalit Bisht, JBIT College of Pharmacy, Dehradun, India

² Ranjana Sonkar, JBIT College of Pharmacy, Dehradun, India

³ Principal, JBIT College of Pharmacy, Dehradun, India

Correspondence should be addressed to Arun Kumar Maurya ³

Arunmedi4u@gmail.com

DOI: 10.47750/pnr.2023.14.03.336

Abstract

Illness bring about reduced cell creation of defensive mixtures prompting expanded harm to cell films; unavoidably, harm layers which lessens cell capacity to fix harmed tissue. Free extreme age has been embroiled in the aggravation of wound region 99 A medication which hinders lipid peroxidation is accepted to expand the reasonability of cells by working on the dissemination, forestalling cell harm, advancing DNA union, and so on and thus advancing the injury mending. Jc oil contains sugars, flavonoids, tannins, and triterpenoids, proteins a polysaccharides and unstable oil. Jc oil contains juniperin, which is tannins answerable for anthelmintic movement. Flavonoids have been reported to have intense cell reinforcement and free extremist rummaging impact, which is accepted to be one of most significant mixtures of wound mending. Subsequently, the upgraded wound mending might be free extremist searching activity of JC, and improved degree of cell reinforcement proteins in the granuloma tissues. Better collagenation seen affected by this JC might be a direct result of improved98. Anyway further investigations are expected to affirm the constituents liable for wound recuperating action. The medication has a critical anthelmintic movement which might be because of both impeding energy digestion and parallelizing the worm. This might be because of tannins. The oil of Juniperus communis linn. was fundamentally powerful in extraction, entry point and dead space wound models. This is additionally affirmed by histopathological studies. The oil has likewise strong hepatoprotective action. The injury recuperating and hepatoprotective movement might be because of presence of dynamic standards like tannins and flavonoids. Nonetheless, further investigations are expected to affirm the constituents answerable for the movement. To assess Juniperus communis linn (JC) oil for anthelmintic, injury recuperating and hepatoprotective exercises in rodents.

Keywords- Juniperus communis linn, anthelmintic, injury recuperating and hepatoprotective movement.

Introduction- Anthelmintic medications are utilized to kill or diminish the quantity of helminthic parasites in the digestive system or tissues of the body. These parasites share numerous biochemical and physiological cycles for all intents and purpose with their human hosts¹. Butea frondasa is a natural medication which is utilized as anthelmintic having palasonin as its dynamic constituent². Natural medications were viewed as invaluable contrasted with engineered and semi manufactured because of its less unfavorable impacts, for example, gastric irritation³. All constant contaminations are brought about by different types of parasitic helminthes. It is assessed that over a portion of the populace might be tainted with gastrointestinal helminthes yet there are inconspicuous contrasts that are starting to respect pharmacological investigation. The skin has been depicted as the biggest organ in the body. Wound has been characterized as the interruption of physical

or practical congruity of living tissue⁴ because of physical, compound, microbial or electrical affront. Wound recuperating is the body's regular course of recovering dermal and epidermal tissue. The mending overflow is enacted when platelets interact with presented collagen prompting platelet collection and arrival of coagulating factors bringing about testimony of fibrin clump at the site of injury. Wound recuperating and fix can happen likewise at whatever point the entire body structure is satisfactory for endurance and working of the particular cells of organ parenchyma. The quickness of wound mending depends to an impressive degree on the withdrawal that starts a couple of days after injury and ceaseless for a few weeks. Ordinarily, the cycle and periods of mending are practically comparable in all warm blooded creatures and progress of recuperating follows an ideal rate which might contrast from species to species and even from one tissue to another in the equivalent species. Curiously examination has uncovered different variables which advance injury recuperating viz. oxygen tension, ascorbic acid⁸, nutrient A⁹, vitamin D, zinc⁹ and there are sure factors which hinder the collagen union and connective tissue fix like, corticosteroids, medroxy progesterone, methyl rednisolone¹⁰ and so on. Coherently, factors which advance or alter any of these occasions upgrade wound recuperating. Wound mending addresses a coordinated work to restore tissue respectability following injury and includes an intricate request of cell and biochemical occasions. The liver is a fundamental organ of principal significance engaged with the support of metabolic elements of central significance associated with the upkeep of digestion capabilities and detoxification from the endogenous and exogenous difficulties, as xenobiotics, drugs, viral contaminations and persistent alcoholism.

Carbon tetrachloride applies its harming impact on the liver cells by change to free extremists. for example transformation of CCl₄ to CCl₃ by the course of dehalogenation of carbon tetrachloride. This dehalogenation is interceded by cytochrome P450, this course of creation of free extremists setting the chain response of lipid peroxidation. This lipid peroxidation increments films penetrability which eventually causes cell ageing. The *Juniperus communis* Linn.(JC) plant is anthelmintic, alexipharmic, diuretic, anti-diarrhoeal, emmenagogue, abortifacient, wound mending, tonic, hepatic, great for ear infection, toothache, heaps, cooling to the mind and cutaneous problems. It is valuable in stomach torments, strangury, sicknesses of spleen and midsection, ascites, growths, heaps, bronchitis, acid reflux, clogging, vaginal release. It has carminative, energizer, diuretic, and sudorific properties. The study of writing uncovers that *Juniperus communis* Linn. is accounted for antimicrobial¹⁴, diuretic¹⁵, antibacterial and antifungal¹⁶ and antioxidant activity . The current review is plan to demonstrate logically the action of JC oil on anthelmintic, injury mending and hepatoprotection.

MATERIAL AND METHODS-

ANTHELMINTIC STUDY:

The current study was led to recognize the anthelmintic action of *Juniperus communis* linn. oil. what's more, the system by which it acts. The in vitro impact of this oil on glucose take-up action of *A.galli* was concentrated on taking Piperazine citrate as a source of perspective medication as influencing this large number of parameters is known. Furthermore, mass motility investigations of *Pheretima posthuma* are likewise finished to distinguish whether these concentrates significantly affect engine activity of the *Pheretima posthuma*. Glucose Uptake Studies:

Materials: *Ascaridia galli* ,Tyrode solution, *Juniperus communis* linn. oil Piperazine citrate Glucose estimation kit (Span Diagnostics)

Method: *A.galli* worms were hatched in two unique fixations (100 g/ml and 300 g/ml) of petrol ether and methanol concentrates of leaves of *Mentha arvensis* Linn. what's more, mebendazole (100 g/ml and 300 g/ml) and medication free Tyrode medium (control) for 20 hours. Assessment of glucose is finished according to Trinder's method.⁸⁴

The glucose assessment was finished by utilizing glucose oxidase/peroxidase (GOD/Unit) technique utilizing a standard pack got from Range Diagnostics.

The items in the unit are compound reagent, support reagent, glucose standard (100mg/dl).

Table No. 3 Glucose uptake studies

| Reagent | Blank (B) | Standard (S) | Test (T) |
|-------------------------|-----------|--------------|----------|
| Serum / Plasma | -- | -- | 0.02 ml |
| Glucose standard 100mg% | -- | 0.02 ml | -- |
| Working glucose reagent | 1.5 ml | 1.5 ml | 1.5 ml |
| Purified water | 1.5 ml | 1.5 ml | 1.5 ml |

Result-

HEPATOTOXIITY :


Grown-up rodents of either sex weighing 150-200g were partitioned into four gatherings each comprising of 6 creatures. Bunch I: Fluid paraffin(3ml/kg) s.c. furthermore, filled in as control. Bunch II: CCL4 + Fluid paraffin(1:2) V/V, 1 ml of CCl4 per kg s.c. filled in as unhealthy control to prompt hepatic harm 24h before beginning of examination. Bunch III: CCL4 + Fluid paraffin+ Silymarin(100mg/kg) p.o. day to day. filled in as standard for 8 days. Bunch IV: CCL4 + Fluid paraffin+ JC (300mg/kg) p.o. day to day for 8 days filled in as test. Blood removed on eighth day from retro-orbital plexus of rodents. Serum was isolated from blood of each rodent by centrifugation for assessment of biochemical boundaries. Rodents were forfeited and liver quickly extracted following penance and put in 10% formalin arrangement sequentially separated and minutely analyzed in the wake of staining with hematoxylin and eosin.

Biochemical Examination :

Assessment of SGPT and SGOT:

Serum glutamate pyruvate transaminase movement as a record of hepatic harm has been generally utilized. Human serum contains a few transaminases of which aspartate transaminase (AST) otherwise called glutamate oxaloacetate transaminase (GOT) and alanine transaminase (ALT) otherwise called glutamate pyruvate transaminase (GPT) are of symptomatic importance. Raised degrees of serum transaminase have been seen in hepatic injury and are generally utilized both clinically and in trial model to uncover the degree of hepatic injury. They are valuable for finding of intense viral hepatitis, liver harm because of medications or synthetic substances and for the separation of these circumstances from obstructive jaundice. Consequently the limit of various concentrates to decrease the raised serum transaminase levels in carbon tetrachloride prompted intense

hepatic disappointment model has been utilized as a biochemical boundary to uncover its hepatoprotective limit.



BIOCHEMICAL PARAMETERS-

Assessment of Serum Glutamate Pyruvate Transaminase (SGPT). 90

SGPT or ALT is situated in the cytosol of the liver cell. During liver cell irritation, they are delivered into course because of expanded porousness of cell layer separate of liver cells. Subsequently, assurance of SGPT as record of the degree of liver harm.

Demonstrative reagent unit was utilized for assurance of SGPT likewise called as "Alanine amino transaminase" (ALT) movement by strategy for Reitman and Frankel. Guideline: SGPT (ALT) Catalyzes the accompanying response: Ketoglutarate + L-Alanine → L-glutamate + pyruvate Pyruvate so shaped is combined with 2,4-Dinitro phenyl hydrazine (2, 4 DNPH) to give the relating hydrazone, which gives earthy colored variety in antacid medium and this can be estimated colorimetrically.

Discussion-

The presence of glucose in the medium, it is hard to survey the impact of the medication straightforwardly on the glucose digestion or glycogen exhaustion might result ensuing to the restraint of glucose take-up and keep up with the typical physiological capability of the parasite. To preclude such a chance, the impact of MBZ and JC on the tissue glycogen level was concentrated on in glucose free medium. Yet, MBZ and the JC neglected to change the glycogen level of *A.galli* without any glucose recommending that MBZ and JC actuated glycogen consumption was because of stamped restraint of glucose take-up which affirms that the medication acts through hindering the energy digestion as in the event that palasonin, a functioning constituent from *Butea frondosa*.⁵ In addition, there is likewise huge abatement in mass motility which shows the movement of medication is likewise by deadening the worm. Phytochemical examination of JC exhibits the presence of sugars, triterpenoids, glycosides and flavonoids. The oil has additionally strong hepatoprotective action. The injury recuperating and hepatoprotective action might be because of presence of dynamic standards like tannins and flavonoids. Nonetheless, further investigations are expected to affirm the constituents answerable for the movement.

Table- Impact of JC oil on different biochemical boundaries

| Groups | BIOCHEMICAL PARAMETERS | | | | |
|------------------------|------------------------|---------------|---------------|----------------|-------------------------|
| | SGPT (IU/L) | SGOT (IU/L) | SALP (IU/L) | TP (g/dl) | Serum Bilirubin (mg/dl) |
| Control | 76.00+6.154 | 166.2+6.306 | 177.7+3.432 | 6.617+0.098 | 0.73+0.024 |
| CCl₄ | 763.3+62.94 | 1238.0+52.68 | 447.2+5.745 | 3.473+0.077 | 2.46+0.087 |
| Silymarin | 243.0+12.43** | 275.5+7.037 | 244.0+10.82** | 6.085+0.1033* | 1.475+0.097** |
| JC | 293.7+22.38** | 591.8+69.72** | 350.2+9.803** | 4.687+0.1928** | 2.458+0.105** |

Effect of JC and Silymarin (standard) on serum levels of SGOT (IU/L), SGPT (IU/L),SALP(IU/L),TP(g/dl) and Bilirubin (mg/dl) level of hepatic tissue during Carbon tetrachloride induced hepatotoxicity in rats. Values are mean \pm S.D. (n = 6). **p < 0.01, *p < 0.05, respectively.

References

1. Betram G. Katzung. "Basics and Clinical Pharmacology", 8th edition. LangeMeical Books, Mcgraw Hill Medical Publishing Division, 2002:903-20.
2. Rang HP, Dale MM, Ritter JM, Moore PK. Anthelmintic drugs, 6th edition,2008Churchill livingstone; Elsevier sites pub: 712-17.
3. Nayak BS, Raju SS, Ramsuhag A. "Investigation of wound healing activity of Lanata camara linn. in Sprague dawley rats using burn wound model". Inter J Appl Res Nat Prod 2008; I: 15-19
4. Sundari PN, Wlifred G, Ramakrishna B.Dose oxidative protein damage play a role in hepathogenesis of carbon tetrachloride- induced liver injury in the rats. Biochem Biophysica Acta 1997; 1362: 169-76.
5. Jay S. Keystone and John K. Murdoch, James W. Tracy and Leslie T. Webster.
6. R.K. Kaushik, J.C. Katiyar, A.B. Sen. Studies on the mode of action of anthelmintics with Ascaridia galli in the test parasite. Indian J Med Res 62; 1367- 75
7. Kataranovski M, Magic Z, Pejnovic N. Early inflammatory cytokine and acute phase protein response under the stress of thermal injury in rats. Physiol Res.1999;48:473-482.
8. Fleming T, Deutsch M, Wyble C.PDR for Herbal Medicines.2nd New Jersey: Medical Economics Company Inc.2000:497,593,690,719,757,821.
9. Thomas Fleming.PDR for herbal medicines.2000: 2:419-420.
10. Asima Chattarjee. The Treatise on Indian medicinal plants. Satyesh ChandraPrakasn. p. 163-165. (Vol 3).
11. Madhan B, Murlidharan C, Jayakumar R.Study on the stabilizaton of Collagen with vegetable tannins in the presence of acrylic polymers. Biomaterials. 2002 :23 (14) : 2841 -2847.
12. Stevenson PC, Simmonds MS, Sampson J, Houghton PJ, Grice P. Wound healing activity of acylated iridoid glycosides from Scrophularia nodosa. Phytoter Res.2002;16(1):33-35.
13. Gupta M, Asunder UK, Vamsi ML, Sivakumar T, Kanddar CC. Antisteroidogenicactivity of the two Indian medicinal plants in mice. J Ethanopharmacol 2004;90:21-5.